



Active Constituents:

ABAMECTIN 5 mg/mL, TRICLABENDAZOLE 300 mg/mL

Product description

AVOMEK PLUS is a convenient pour-on treatment that not only controls worms and external parasites but has the added benefit of controlling all three stages of liver fluke.

Product benefits

Pour-On

- ✓ Convenient
- ✓ Simple and safe
- ✓ Rainfast

Internal and External Parasite Control

- ✓ Worms
- ✓ Lice
- ✓ Cattle Tick (aids in control)

Liver Fluke - Eliminates all 3 stages

- ✓ Early immature
- ✓ Immature
- ✓ Adult

Liver Fluke Disease

In cattle, liver fluke infection has been shown to result in reduced fertility, milk yield, feed conversion, carcass yield and growth rates. Infections are usually subclinical and 8-9% reductions in weight gain can occur with no clinical signs of disease¹.

The most severe damage caused by liver fluke infection in livestock is the result of the early immature and immature stages migrating through the liver tissue on their way to the bile ducts where they subsequently reside as adults.

Strategic treatment of all stock with a product that kills all three stages of liver fluke has the dual benefit of:

1. Removing resident liver fluke infection.
2. Preventing pasture contamination with fluke eggs at a time when fluke development and snail activity in the environment is about to recommence. This reduces fluke infection in livestock.

The use of a flukicide which contains triclabendazole*, the only active that kills all three stages of liver fluke, is recommended best practice on farms with a known fluke problem.

AVOMEK PLUS contains triclabendazole which effectively kills all three stages of liver fluke. Triclabendazole reduces fluke burdens and pasture contamination, which improves the productivity of infected livestock.

Cattle Indications

When used as directed, AVOMEK PLUS treats and controls the following parasites of cattle:

Gastrointestinal roundworms:

- Barber's Pole Worm (*Haemonchus placei*)
- Small Brown Stomach Worm (*Ostertagia ostertagi*) – including inhibited stages
- Small Intestinal Worm (*Cooperia oncophora*, *Cooperia punctata*)
- Stomach Hair Worm (*Trichostrongylus axei*)
- Hookworm (*Bunostomum phlebotomum*)
- Large Bowel Worm (*Oesophagostomum radiatum*)
- Whipworm (*Trichuris* spp.)
- Thin Necked Intestinal Worm (*Nematodirus spathiger*)
- Intestinal Threadworm (*Strongyloides papillosus*)

Lungworm (*Dictyocaulus viviparus*)

Sucking and Biting Lice

- Long-nosed sucking louse (*Linognathus vituli*)
- Short-nosed sucking louse (*Haematopinus eurysternus*)
- Little sucking louse (*Solenopotes capillatus*)

Biting louse (*Bovicola bovis*)

Liver Fluke (*Fasciola hepatica*)

- Early immature – 4 week old larvae
- Immature – 6 week old larvae
- Adult – 12 weeks and over

Also aids in the control of:

Cattle Tick (*Boophilus microplus*) – Not recommended for use in strategic dipping programs for cattle tick control.

Buffalo Fly (*Haematobia irritans exigua*)



Dose rate

1 mL/10 kg bodyweight. Delivering 0.5 mg abamectin and 30 mg triclabendazole per kg bodyweight.

It is applied along the back line of the animal in a strip starting between the shoulder blades. AVOMEK PLUS should not be applied to calves under 50 kg bodyweight.

Live weight (kg)	Dose Volume (mL)	1 Litre Pack treats	2.5 Litre Pack treats	5 Litre Pack treats	15 Litre Pack treats
51-75	7.5	133	333	666	2000
76-100	10	100	250	500	1500
101-125	12.5	80	200	400	1200
126-150	15	66	166	333	1000
151-175	17.5	57	142	285	857
176-200	20	50	125	250	750
201-250	25	40	100	200	600
251-300	30	33	83	166	500
301-350	35	28	71	142	428
351-400	40	25	62	125	375
401-450	45	22	55	111	333
451-500	50	20	50	100	300
501-550	55	18	45	90	272
551-600	60	16	41	83	250
601-650	65	15	38	76	230

Withholding Periods

MEAT: DO NOT USE less than 49 days before slaughter for human consumption.

MILK: DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption.

EXPORT SLAUGHTER INTERVAL (ESI): 140 days.

Storage

Store below 30°C (room temperature).

Pack sizes

Available in 1 L, 2.5 L and 5 L packs, and 15 L drum.

Both the 2.5 L and 5 L pack sizes are packaged in a convenient backpack for easy use.



*See product label for full claim details and directions for use. 1. Hope-Cawdrey et al. (1997) Production effects on liver fluke in cattle Part I. the effects of infection on liveweight gain, feed intake and food conversion efficiency in beef cattle, British Vet], 133 (2): 145-159. Boehringer Ingelheim Animal Health Australia Pty. Ltd., Level 1, 78 Waterloo Road, North Ryde, NSW 2113 Australia. ABN 53 071 187 285. *AVOMEK is a registered trademark of the Boehringer Ingelheim Group. All rights reserved. AUS/AVPL-181000

