



Eprinex®

Pour-On for Beef and Dairy Cattle



Active Constituent:
5 mg/mL EPRINOMECTIN

Product description

For the treatment and control of internal and external parasites of beef and dairy cattle and internal parasites of deer.

Product benefits

- ✓ The convenience of a broad-spectrum pour-on
- ✓ Sustained activity to kill more species of worms for longer than any other pour-on*
- ✓ Nil milk, nil meat withholding periods and nil ESI
- ✓ Completely weatherproof – not just rainfast

Cattle Indications

When used as directed EPRINEX treats and controls susceptible strains of the following parasites of cattle:

Gastrointestinal roundworms

For the control of adult and immature:

- Ostertagia* spp.
- Ostertagia ostertagi* (small brown stomach worm) (including inhibited L4)
- O.lyrata* (brown stomach worm) (adults only)
- Haemonchus placei* (barber's pole worm)
- Trichostrongylus* spp.
- Trichostrongylus axei* (stomach hair worm)
- T. colubriformis* (intestinal hair worm)
- Cooperia* spp. (small intestinal worm) (including inhibited L4)
- C. oncophora*
- C. pectinata*
- C. punctata*
- C. surnabada*
- Oesophagostomum* spp.

- Oesophagostomum radiatum* (nodule worm)
- Nematodirus helvetianus* (thin necked intestinal worm)
- Bunostomum phlebotomum* (hookworm)
- Trichuris* spp. (whip worm) (adults only)

Lungworm

Adult and immature *Dictyocaulus viviparus*

Sucking And Biting Lice

- Linognathus vituli*
- Haematopinus eurysternus*
- Bovicola (Damalinia) bovis*
- Solenopotes capillatus*

Mites

- Chorioptes bovis*
- Sarcoptes scabiei* var *bovis*

Buffalo Fly

Haematobia irritans exigua

Aids in control of:

Cattle Tick *Boophilus microplus*

Days of protection offered against reinfection

	<i>Ostertagia</i> spp. <i>O. ostertagi</i> <i>O. lyrata</i> <i>O. leptospicularis</i> <i>Cooperia</i> spp. <i>C. oncophora</i> <i>C. pectinata</i> <i>C. surnabada</i> <i>Trichostrongylus</i> spp. <i>T. axei</i> <i>T. colubriformis</i> <i>Nematodirus helvetianus</i> <i>Oesophagostomum radiatum</i> <i>Haemonchus placei</i> <i>Bunostomum phlebotomum</i> <i>Dictyocaulus viviparus</i>																Days of Persistent Activity
EPRINEX® Pour-On	28*	28*	28*	28*	28*	28*	28*	28*	21	21	21	28*	28*	21	NC	28	
Cyductin® Pour-On	NC	42	42	NC	NC	NC	NC	NC	NC	28	NC	21	42	28	42	42	
Dectomax® Pour-On	NC	35*	NC	NC	35*	21*	NC	NC	NC	35*	NC	NC	21*	35*	21*	28*	

NC = NO registered label claim for persistent activity. Persistent activity is the length of time a product continues to kill incoming parasite larvae after administration. *Up to the periods shown. See product label for full claim details.





Eprinex

Pour-On for Beef and Dairy Cattle



Directions for use

Dosage and Administration

A representative sample of animals should be weighed with either scales or a weighband. Dose the mob according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead. The dose rate is 1 mL for each 10 kg of liveweight.

Live Weight (kg)	Dose Volume (mL)	Number of doses per pack size			
		1 L	2.5 L	5 L	20 L
Up to 50	5	200	500	1000	4000
51 - 75	7.5	133	333	666	2666
76 - 100	10	100	250	500	2000
101 - 125	12.5	80	200	400	1600
126 - 150	15	66	166	333	1333
151 - 175	17.5	57	142	285	1142
176 - 200	20	50	125	250	1000
201 - 250	25	40	100	200	800
251 - 300	30	33	83	166	666
301 - 350	35	28	71	142	571
351 - 400	40	25	62	125	500
401 - 450	45	22	55	111	444
451 - 500	50	20	50	100	400
501 - 550	55	18	45	90	363
551 - 600	60	16	41	83	333
601 - 650	65	15	38	76	307

Heavier mature bulls should receive an additional 5 mL for each 50 kg over 650 kg.

Cattle Withholding Periods

MEAT: NIL

MILK: NIL

EXPORT SLAUGHTER INTERVAL (ESI): NIL

Rainfast

EPRINEX is not just rainfast, it is fully weatherproof. This gives the flexibility to use at any time, in any weather conditions. Rainfall before or after treatment will not affect the efficacy of EPRINEX Pour-On.

Storage

Store below 30°C (room temperature). Store bottle or pack in carton to protect from light.

Pack sizes

Available in 1 Litre, 2.5 Litre, 5 Litre and 20 Litre packs. Both the 2.5 L and 5 L pack sizes are packaged in convenient, collapsible backpacks for easy use. The 1 L pack sizes feature the flexible 'Magicap' dispenser that eliminates the need for a drench gun.



*See product label for full claim details and directions for use. *Up to the periods shown in Table 1. Boehringer Ingelheim Animal Health Australia Pty. Ltd., Level 1, 78 Waterloo Road, North Ryde, NSW 2113 Australia. ABN 53 071 187 285. EPRINEX® is a registered trademark of the Boehringer Ingelheim Group. *Cydectin is a registered trademark of Virbac. *Dectomax is a registered trademark of Zoetis. All rights reserved. AUS/IVEP-181008