

FENNEC

POUR-ON FOR SHEEP



Australia's first dual active combination pour-on lousicide for sheep.

Product description

For the treatment and control of susceptible strains of lice (*Bovicola ovis*) on sheep, including synthetic pyrethroid resistant lice when applied up to 7 days after shearing.

Sheep will be protected against re-infestation with lice for 5 weeks after treatment when applied within 24 hours of shearing.

For the treatment and control of abamectin susceptible gastrointestinal nematodes of sheep.

Fennec™ Pour-On for Sheep contains imidacloprid, a member of the neonicotinoid family of chemicals, and abamectin, a member of the macrocyclic lactone family of chemicals. Resistance may develop to any chemical.

Active Constituents:

35 mg/mL Imidacloprid

4 mg/mL Abamectin

Solvent: 300 mg/mL N-methyl-2-pyrrolidone

Product benefits

- ✓ Australia's first combination pour-on for lice
- ✓ 5 weeks protection from re-infestation with lice when treated when treated within 24 hours of shearing
- ✓ No known resistance in lice to either active
- ✓ Two modes of action deployed simultaneously
- ✓ Convenient off-shears pour on application

Contraindications

DO NOT USE on unshorn sheep, or sheep more than 7 days after shearing.

DO NOT USE in ewes with unshorn lambs at foot.

This product should not be used on lambs under 6 weeks of age or less than 10 kg bodyweight.

This product should not be used on severely sick, debilitated or dying animals.

This product should not be used on sheep which have not been cleanly shorn for any reason e.g. Mycotic Dermatitis.

Sheep Indications

Fennec™ Pour-On for Sheep is effective for the treatment and control of the following abamectin sensitive adult and immature gastrointestinal nematodes:

Small Brown Stomach Worm (*Teladorsagia circumcincta*)

Black Scour Worm (*Trichostrongylus* spp.)

Stomach Hair Worm (*Trichostrongylus axei*)

Nodule Worm (*Oesophagostomum columbianum*)

Large Bowel Worm (*Oesophagostomum venulosum*)

Restrictions

DO NOT USE on ewes which are producing, or may in the future produce milk that may be used or processed for human consumption.



FENNEC

POUR-ON FOR SHEEP



Dose rate

10 mL for each 10 kg body weight. Sheep should be weighed before treatment.

Apply the entire dose to sheep under 30 kg as a single broadband stripe from poll of the head to the base of the tail.

For sheep over 30 kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the base of the tail. The stripes should be no further than 10 cm apart. Refer to diagrams below. The following table may be used as a guide:

Bodyweight (kg)	Dose Volume (mL)	20 L Treats	Method	Apply in the pattern below
10 to 12.5	12.5	1600	Single stripe	
12.6 to 15	15	1333		
15.1 to 20	20	1000		
20.1 to 25	25	800		
25.1 to 30	30	666		
30.1 to 40	40	500	Double stripe	
40.1 to 50	50	400		
50.1 to 60	60	333		
60.1 to 70	70	285		
70.1 to 80	80	250		
80.1 to 90	90	222		

Animals heavier than 90 kg to be dosed at 10 mL per 10 kg. A representative sample of animals should be weighed before treatment. Dose the mob according to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Withholding Periods

MEAT: DO NOT USE less than 63 days before slaughter for human consumption.

MILK: DO NOT USE on ewes which are producing, or may in the future produce milk that may be used or processed for human consumption.

WOOL: DO NOT USE less than 6 months before shearing or fibre collection.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 84 days before slaughter for export. Before using this product, confirm the current ESI from Boehringer Ingelheim on 1800-808-691 or the APVMA website (www.apvma.gov.au/residues).

Storage

Store below 30°C (Room Temperature). Protect from light.

Pack size

20 L drum.



*See product label for full claim details and directions for use. Boehringer Ingelheim Animal Health Australia Pty. Ltd., Level 1, 78 Waterloo Road, North Ryde NSW 2113 Australia. ABN 53 071 187 285. *Fennec is a trademark of the Boehringer Ingelheim Group. AUS/GENS-181006