

ivomec®

(ivermectin) **Pour-On** for Cattle



Active Constituent:

5 mg/mL IVERMECTIN

Product description

For the treatment and control of ivermectin-sensitive gastrointestinal roundworms, lungworm, eyeworm, sucking and biting lice, sarcoptic and chorioptic mange mites, buffalo fly and cattle tick in cattle, and for the treatment and control of gastrointestinal roundworms and lungworm of red deer.

Product benefits

- ✓ Controls worms, lice, cattle tick and buffalo fly
- ✓ Buffalo fly control – up to 21 days
- ✓ Long cattle tick control – 28 days[#]
- ✓ Long lice protection – up to 56 days[#]
- ✓ Convenience of a pour-on application
- ✓ Wide margin of safety when used as directed
- ✓ Rainfast

Persistent activity

At the recommended dose of 1 mL per 10 kg, IVOMEc Pour-On controls:

- ✓ *Trichostrongylus axei* and *Cooperia* spp. for up to 14 days after treatment;
- ✓ *Ostertagia* spp. and *Oesophagostomum radiatum* for up to 21 days after treatment;
- ✓ *Dictyocaulus viviparus* (Lungworm) for up to 28 days after treatment;
- ✓ *Haematobia irritans exigua* (Buffalo Fly) for up to 21 days after treatment;
- ✓ Lice (*Bovicola bovis* and *Linognathus vituli*) for up to 56 days after treatment;
- ✓ Prevents the appearance of viable adult cattle ticks (*Boophilus microplus*) on cattle for 28 days after treatment.

Cattle Indications

When used as directed, IVOMEc Pour-On treats and controls the following parasites of cattle:

Gastrointestinal roundworms:

Adult and immature (L₄):

- Small brown stomach worm (*Ostertagia ostertagi*) (including inhibited stages)
- Barber's pole worm (*Haemonchus placei*)
- Stomach hair worm (*Trichostrongylus axei*)
- Intestinal hair worm (*T. colubriformis*)
- Small intestinal worm (*Cooperia* spp.)
- Nodule worm (*Oesophagostomum radiatum*)

Adult stages of:

- Cooperia oncophora*
- C. punctata*
- Oesophagostomum venulosum*
- Whipworm (*Trichuris* spp.)
- Intestinal threadworm (*Strongyloides papillosus*)

Immature stages of:

- Thin necked intestinal worm (L₄) (*Nematodirus* spp.)

Lungworm:

- Dictyocaulus viviparus* (adult and immature)

Eyeworm

- Thelazia* spp. (adult stages)

Sucking and Biting Lice

- Linognathus vituli*
- Haematopinus eurysternus*
- Bovicola (Damalinia) bovis*
- Solenopotes capillatus*

Buffalo Fly

- Haematobia irritans exigua*

Mites

- Chorioptes bovis*
- Sarcoptes scabiei* var. *bovis*

Cattle Tick

- Boophilus microplus* (including organophosphate-, synthetic pyrethroid- and amidine- resistant strains)

For indications for deer, see label for full details and directions of use.

ivomec®

(ivermectin) **Pour-On** for Cattle



Dose rate

1 mL for each 10 kg liveweight.

Live Weight (kg)	Dose Volume (mL)	Number of doses per pack size			
		1 L pack	2.5 L pack	5 L pack	20 L pack
50 - 100	10	100	250	500	2000
101 - 150	15	66	166	333	1333
151 - 200	20	50	125	250	1000
201 - 250	25	40	100	200	800
251 - 300	30	33	83	166	666
301 - 350	35	28	71	142	571
351 - 400	40	25	62	125	500
401 - 450	45	22	55	111	444
451 - 500	50	20	50	100	400
501 - 550	55	18	45	90	363
551 - 600	60	16	41	83	333
601 - 650	65	15	38	76	307

Heavier animals (for example mature bulls) should receive an additional 5 mL for each 50 kg over 650 kg.

Withholding Periods

Meat: DO NOT APPLY to cattle less than 42 days before slaughter for human consumption.

DO NOT APPLY to deer less than 35 days before slaughter for human consumption.

MILK: When milk or milk products are to be used for human consumption do not administer to cows within 28 days prior to calving or during lactation.

EXPORT SLAUGHTER INTERVAL (ESI): 42 days

Storage

Store below 30°C (room temperature).
Store bottle in carton to protect from light.

Pack sizes

Available in 1 Litre, 2.5 Litre, 5 Litre & 20 Litre packs.
The 20 L pack size is packaged in a robust drum for the larger producer.
Both the 2.5 L and 5 L pack sizes are packaged in convenient, collapsible backpacks for easy use.
The 1 L pack size features the flexible 'Magicap' dispenser which eliminates the need for a drench gun.



#See product label for full claim details and directions for use. Boehringer Ingelheim Animal Health Australia Pty. Ltd., Level 1, 78 Waterloo Road, North Ryde, NSW 2113 Australia. ABN 53 071 187 285. *IVOMECE is a registered trademark of the Boehringer Ingelheim Group. All rights reserved. AU-CAT-0044-2024