

TRIGUARD®

Triple Combination Drench for Sheep



Active Constituents:

1.0 g/L ABAMECTIN, 22.7 g/L OXFENDAZOLE,
33.9 g/L LEVAMISOLE (as levamisole hydrochloride),
0.5 g/L SELENIUM (as sodium selenate),
2.2 g/L COBALT (as cobalt disodium EDTA)

Product description

For the control and treatment of internal parasites of sheep including those with single or dual resistance to macrocyclic lactones (including moxidectin resistant strains), benzimidazoles or imidazothiazoles (levamisole / morantel).

Also provides a selenium and cobalt supplement for sheep raised in selenium and/or cobalt deficient areas.

Product benefits

- ✓ Combined power of 3 actives¹
- ✓ No on-farm mixing
- ✓ Kills worms resistant to white, clear, or ML drenches
- ✓ Can significantly delay the development of resistance
- ✓ Stable formulation
- ✓ High potency with broad-spectrum activity
- ✓ Proven in the field

Broad-spectrum activity against resistant worms

With high potency triple-combination protection, TRIGUARD can deliver highly effective, broad-spectrum worm control even where high levels of resistance are present.

TRIGUARD is active against sheep roundworms that have single or dual resistance to the MLs (including moxidectin), benzimidazole or imidazothiazole (levamisole) families of chemicals.

Sheep Indications

At the recommended dosage level, TRIGUARD provides treatment and control of the following sheep roundworms, sensitive to at least one of the three antiparasitic components of the product:

Adult and immature gastrointestinal roundworms

Barber's Pole Worm (*Haemonchus contortus*)[^]
Small Brown Stomach Worm (*Teladorsagia [Ostertagia] circumcincta*)[^]
Stomach Hair Worm (*Trichostrongylus axei*)
Black Scour Worm (*Trichostrongylus* spp.)
Small Intestinal Worm (*Cooperia* spp.)[^]
Thin Necked Intestinal Worm (*Nematodirus* spp.)
Hookworm (*Bunostomum trigonocephalum*)
Intestinal Threadworm (*Strongyloides papillosus*)
Large Bowel Worm (*Oesophagostomum venulosum*)
Large Mouthed Bowel Worm (*Chabertia ovina*)
Nodule Worm (*Oesophagostomum columbianum*)
Whipworm (*Trichuris ovis*)

[^]including inhibited fourth stage larvae.

Lungworm (*Dictyocaulus filaria*)

Also for the control of:

Nasal Bot (*Oestrus ovis*) parasitic larval stages only.

Itch Mite (*Psorergates ovis*)



TRIGUARD[®]

Triple Combination Drench for Sheep



Dose rate

1 mL / 5 kg body weight, which provides 0.2 mg abamectin, 4.5 mg oxfendazole, 6.8 mg levamisole (8 mg levamisole hydrochloride), 0.1 mg selenium and 0.44 mg cobalt per kg bodyweight.

| Live weight (kg) | Dose (mL) | 5 L pack treats | 20 L pack treats |
|------------------|-----------|-----------------|------------------|
| 16-20 | 4 | 1250 | 5000 |
| 21-25 | 5 | 1000 | 4000 |
| 26-30 | 6 | 833 | 3332 |
| 31-35 | 7 | 714 | 2856 |
| 36-40 | 8 | 625 | 2500 |
| 41-50 | 10 | 500 | 2000 |
| 51-60 | 12 | 416 | 1666 |
| 61-70 | 14 | 357 | 1428 |
| 71-80 | 16 | 312 | 1250 |

Withholding Periods

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

MILK: DO NOT USE in ewes, which are producing or may in the future produce milk or milk products for human consumption.

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT use less than 28 days before slaughter for export.

Storage

Store below 30°C (room temperature) away from light.

Pack sizes

5 L backpack and 20 L drum.



See product label for full claim details and directions for use. 1. Data on file Triguard trial data summary 2009-2012. Boehringer Ingelheim Animal Health Australia Pty. Ltd., Level 1, 78 Waterloo Road, North Ryde, NSW 2113 Australia. ABN 53 071 187 285. TRIGUARD[®] is a registered trademark of the Boehringer Ingelheim Group. All rights reserved. AU-MSP-0158-2024